

Confinement, Turbulence and Diffraction Catastrophes

J.-P. Blaizot^a and M.A. Nowak^b

^a*IPhT, CEA-Saclay, 91191 Gif-sur-Yvette, France*

^b*M. Smoluchowski Institute of Physics, Jagiellonian University, PL-30-059 Cracow, Poland*

Abstract

Many features of the large N_c transition that occurs in the spectral density of Wilson loops as a function of loop area (observed recently in numerical simulations of Yang-Mills theory by Narayanan and Neuberger) can be captured by a simple Burgers equation used to model turbulence. Spectral shock waves that precede this asymptotic limit exhibit universal scaling with N_c , with indices that can be related to Berry indices for diffraction catastrophes.

Key words: Wilson loop, Burgers equation, random matrix

1. Spectral shock waves

Recent lattice studies [1] have provided evidence for a novel universal phenomenon in Yang-Mills theory with a large number of colors N . For small loops, the eigenvalues of the unitary Wilson operator concentrate around the point $z = 1$ in the complex plane, and spread symmetrically on the unit circle towards $z = -1$ as the area of the loop increases. At some critical area, the spectral gap closes, and the corresponding level density is given by a universal function (Pearcey function), whose two arguments exhibit scalings in $N^{1/2}$ and $N^{3/4}$, respectively, apparently independently on the dimensionality of the Yang-Mills theory. For very large loops, the level density goes towards a universal distribution, reflecting disorder, as suggested long ago by Durhuus and Olesen [2]. In a recent paper [3], we suggested a particular mechanism leading to such transition, by noting that in $d = 2$, the spectral evolution of Wilson loops is governed by a complex Burgers equation

$$\partial_A F + F \partial_\theta F = 0, \quad F(\theta, A) = \frac{1}{2} \int_{-\pi}^{\pi} d\alpha \rho(\alpha, A) \cot((\theta - \alpha)/2) \quad (1)$$

where the imaginary part of the resolvent $F(\theta, A)$ gives spectral density ρ . All known properties of the Durhuus and Olesen transition can be easily recovered by analyzing

the solution of Eq. (1), using the methods of complex characteristics, and following the motion of singularities in the complex plane [3]. In this description, the transition corresponds to the collision of two “tsunami-like” shock waves (associated to the edges of the spectrum) evolving symmetrically from $z = 1$ around the unit circle, to merge violently at $z = -1$ at the critical area. For larger areas, the cascade of Fourier modes evolves in such a way that only the lowest mode, corresponding to the size of the system, survives, a phenomenon reminiscent of the inverse turbulent cascade.

2. Non-linear diffusion, universality and colored catastrophes

The complex Burgers equation describes, under certain conditions, the random walk of eigenvalues of large unitary matrices [4]. The role of the time t is played by the area of the loop. Writing the eigenvalues as $z = e^{i\theta}$, the Langevin equation corresponding to the random walk of unitary matrices reads

$$d\theta_i = dB_i + \frac{1}{2} \sum_{i \neq j} \cot \frac{\theta_i - \theta_j}{2} dt \quad (2)$$

where the first term is a Brownian noise and the second is originating from the usual repulsion of eigenvalues. From the Langevin equation, a Smoluchowski-Fokker-Planck equation is obtained using standard techniques, which, after a suitable approximation yields the following equation for ρ [6]

$$\partial_t \rho + \partial_\theta [\rho H[\rho]] = \frac{1}{2} \partial_{\theta\theta} \rho, \quad H[\rho] \equiv \frac{1}{2\pi} \text{P.V.} \int_{-\pi}^{\pi} d\alpha \rho(\alpha, A) \cot \frac{\theta - \alpha}{2}, \quad (3)$$

where P.V. denotes the principal value of the integral. Eq. (3) is equivalent to the viscid Burgers equation, the last term playing the role of a viscosity term. This equation exhibits various scaling with N , which we now discuss.

- *Dyson gas*: We rescale the density, originally normalized to N , as $\rho = N\hat{\rho}$, and the time as $\tau = Nt$. Then, in the large N limit we get [5] (equivalent to Eq. 1)

$$\partial_\tau \hat{\rho} + \partial_\theta [\hat{\rho} H[\hat{\rho}]] = 0. \quad (4)$$

Note that the viscous term has disappeared, since it was dwarfed by the effective spectral viscosity, $v_s = 1/2N^{-1}$. In this regime the evolution proceeds without collisions.

- *Dyson liquid*: There is however another regime, where collisions between the eigenvalues, taken into account by the viscous term, cannot be ignored. This happens in the vicinity of the shock waves, and seems to be responsible for the universal scaling at the edges of the spectrum. To approach this limit, we set $\rho(\theta, t) = N^{\alpha+1} \tilde{\rho}(x, \tau)$ with $x \equiv N^\alpha \theta$ and $\tau \equiv N^{2\alpha} t$, and get

$$\partial_\tau \tilde{\rho}(x, \tau) + N \partial_x (\tilde{\rho} H[\tilde{\rho}]) = \frac{1}{2} \partial_{xx} \tilde{\rho}. \quad (5)$$

This viscid complex Burgers equation allows us to describe the scaling properties of the spectral density in the vicinity of a shock.

¹ This value of the spectral viscosity has also been obtained by Neuberger [9] using different arguments.

This can be seen schematically by applying the Cole-Hopf trick to solve the equation [7]. The resulting spectral distribution can be written in the form

$$\tilde{\rho}(x) = -\frac{1}{N\pi} \text{Im} \partial_x \ln K(x) \quad (6)$$

where $K(x)$ is the convolution of a Gaussian kernel with $K_0(z) = \exp\{-N \int^z \tilde{F}_0(w) dw\}$, where $\tilde{F}_0$ is the resolvent corresponding to the appropriate initial condition for the microscopic spectral density $\tilde{\rho}$. In the vicinity of the shock waves, the resolvent can be approximated by a polynomial of second order (gapped phase) or third order (at the closure of the gap). The resulting spectral densities are therefore related to complex Airy or Pearcey type integrals, with well defined scaling properties.

In fact these scaling properties appear to be universal and manifest themselves in various physical contexts. We mention here an amusing analogy with optics. In geometric optics (vanishing wavelength λ), the rays of light may condense on surfaces of infinite intensity, the so-called caustics. These correspond to stable singularities that are classified, the lowest two corresponding to fold and cusp singularities. At finite wavelength, interference effects enter, and we have to use wave packets to describe the emergence of singularities. Such a wave packet Ψ is generically of the form [8]

$$\Psi(\vec{r}, \lambda) = \frac{1}{\lambda^\beta} \psi\left(\frac{x_1}{\lambda^{\sigma_1}}, \frac{x_2}{\lambda^{\sigma_2}}, \dots\right) \quad (7)$$

For the fold, $\beta = 1/6$ and $\sigma = 2/3$, and Ψ is given by an Airy function, while for the cusp $\beta = 1/4$, $\sigma_1 = 1/2$, $\sigma_2 = 3/4$, and Ψ is given by a Pearcey function [8]. The analogy with these “diffractive catastrophes” goes as follows. The limit of geometrical optics corresponds to the limit $N = \infty$ ($\lambda \sim 1/N$). Rays of light correspond to the (complex) characteristics of the Burgers equation, whose envelopes correspond to the caustics and are associated to singularities of Burgers equation. The universal behavior in the vicinity of these singularities is captured by a Pearcey function, and the particular critical exponents for scaling with “angle” and “area” observed in [1] are identical to the Berry indices for a cusp catastrophe.

This work was supported by Marie Curie TOK Grant MTKD-CT-517186 “Correlations in Complex Systems” (COCOS).

References

- [1] R. Narayanan and H. Neuberger, JHEP 0712 (2007) 066.
- [2] B. Durhuus and P. Olesen, Nucl. Phys. **B184**(1981) 461.
- [3] J.-P. Blaizot and M.A. Nowak, Phys. Rev. Lett. **101** (208) 100102.
- [4] F.J. Dyson, J. Math. Phys. **3** (1962) 1191.
- [5] P. Biane and R. Speicher, Ann. Inst. H. Poincaré PR **37** (2001) 581.
- [6] J.-P. Blaizot and M.A. Nowak, in preparation.
- [7] E. Cépa and D. Lépingle, ESAIM:Probability and Statistics **5** (2001) 203.
- [8] see e.g. M.V. Berry and S. Klein, Proc. Natl. Acad. Sci. USA **93** (1996) 261.
- [9] H. Neuberger, Phys. Lett. **B666** (2008) 106 and *ib.* **B670** (2008) 235.